

Charge fluctuations in a quark-antiquark system

A.Bialas

M.Smoluchowski Institute of Physics
Jagellonian University, Cracow*

February 1, 2008

Abstract

It is pointed out that the recent data on charge fluctuations observed in heavy ion collisions are compatible with production of a system of weakly correlated constituent quarks and antiquarks.

It was recently suggested [1, 2] (see also [3, 5, 4]) that measurements of the quantity

$$D = 4 \frac{\langle \delta Q^2 \rangle}{\langle N_{ch} \rangle} \quad (1)$$

can be used to distinguish the hadron gas in equilibrium from the quark-gluon plasma.

It was then argued in [3] that -after appropriate corrections for resonance production are taken into account- $D \approx 3$ for the hadron gas whereas for the quark-gluon plasma $D \approx 1$.

The preliminary data from CERES [6], NA49 [7] and STAR [8] experiments indicate that the measured value of D is close to that predicted for hadron gas and differs markedly from that expected for QGP, i.e. for a weakly correlated quark-gluon system.

*Address: Reymonta 4, 30-059 Krakow, Poland; e-mail: bialas@th.if.uj.edu.pl;

This rises a question: what kind of structure of the partonic system produced at the early stage of collisions may be compatible with these experimental results? The present note is an attempt to answer this question.

The first observation is that the estimate of the numerator in (1) is rather straightforward. The argument (essentially the repetition of the argument given in [2]) goes as follows. Let us consider a system of several particle species (labelled by i) with charges q_i and multiplicities n_i . Since

$$Q = \sum_i q_i n_i \quad \rightarrow \quad \langle Q \rangle = \sum_i q_i \langle n_i \rangle \quad (2)$$

we obtain

$$\langle \delta Q^2 \rangle \equiv \langle Q^2 \rangle - \langle Q \rangle^2 = \sum_i (q_i)^2 \langle n_i \rangle + \sum_{i,k} c_{ik}^{(2)} \langle n_i \rangle \langle n_k \rangle q_i q_k \quad (3)$$

where $c_{ik}^{(2)}$ are the normalized two-particle correlation functions:

$$c_{ii}^{(2)} = \frac{\langle n_i(n_i - 1) \rangle}{\langle n_i \rangle^2} - 1; \quad c_{ik}^{(2)} = \frac{\langle n_i n_k \rangle}{\langle n_i \rangle \langle n_k \rangle} - 1 \quad \text{if } i \neq k \quad (4)$$

If particles are uncorrelated the second term in (3) vanishes. Moreover, one notes that the first term is the sum of only positive quantities, whereas in the second term cancellations are possible. For example, if all normalized two-particle correlations are identical, one obtains

$$\langle \delta Q^2 \rangle = \sum_i (q_i)^2 \langle n_i \rangle + c^{(2)} \langle Q \rangle^2. \quad (5)$$

and thus the second term disappears when $\langle Q \rangle = 0$.

From now on we shall thus restrict the discussion to the case when the second term can be neglected and thus

$$\langle \delta Q^2 \rangle = \sum_i (q_i)^2 \langle n_i \rangle \quad (6)$$

For pion gas this means

$$\langle \delta Q^2 \rangle = \langle n_+ \rangle + \langle n_- \rangle = \langle N_{ch} \rangle \quad (7)$$

and thus

$$D = 4. \quad (8)$$

For a quark-gluon system we obtain

$$\langle \delta Q^2 \rangle = \frac{1}{9}(4 \langle n_u \rangle + \langle n_d \rangle + 4 \langle n_{\bar{u}} \rangle + \langle n_{\bar{d}} \rangle) \quad (9)$$

(gluons of course do not contribute). In the simplest case when abundances of all quarks are identical we have

$$\langle \delta Q^2 \rangle = \frac{5}{18} \langle N_q \rangle \quad (10)$$

where $\langle N_q \rangle$ is the total number of quarks and antiquarks.

To calculate D it is, however, also necessary to estimate $\langle N_{ch} \rangle$ and this is of course model-dependent. The question is the role of gluons. If they decay into $q - \bar{q}$ pairs before making hadrons, the number of hadrons may be very large. This is what happens in the QGP scenario. And this is the reason of the small value of D obtained in [1, 2]. The argument is based on entropy. Qualitatively: since gluons have a large entropy, they have to produce many particles. Some of them will be charged and thus contribute to the denominator of (1).

If one considers the $q - \bar{q}$ coalescence scenario [9, 10], however, the result is very different. In this scenario gluons are attached to quarks (forming constituent quarks) and thus do not provide important contribution to entropy. Hadrons are created simply by coalescence of the (constituent) $q - \bar{q}$ pairs. Consequently, the total number of hadrons is about 1/2 of the total number of quarks and antiquarks:

$$\langle N_h \rangle = \frac{1}{2} \langle N_q \rangle \quad (11)$$

It is natural to assume that 2/3 of them are charged, so that we obtain

$$\langle N_{ch} \rangle = \frac{2}{3} \langle N_h \rangle = \frac{1}{3} \langle N_q \rangle. \quad (12)$$

and thus

$$D = \frac{10}{3} = 3.333 \quad (13)$$

which is only slightly smaller than (8).

The values of D given by (8) and (13) should still be corrected for resonance production [2]. Since the difference between them is fairly small, however, one should not expect a great difference between the corrected results, either.

We thus conclude that the data on charge fluctuations are not incompatible with the idea that a (quasi)uncorrelated constituent quark system is created before hadronization. This observation strenghtens earlier arguments leading to the coalescence picture of hadronization (see e.g. [9] -[13]).

Two comments are in order.

(i) The present data are given for fairly small rapidity intervals and thus important effects of charge transport between the boundaries of the considered phase-space region may be present [14, 15], destroying the argument of Refs. [1] and [2]. This effect may perhaps also be responsible for the rather large values of D (exceeding 4) observed at SPS energies [6, 7]. Data for larger intervals would be very useful to sort out this question.

(ii) An argument similar to that presented here can be also applied to fluctuations of the net baryon number [1, 4]. The formulae (2) - (6) are valid with the substitution of the charges by baryon numbers. In particular, for the hadron gas we have

$$\langle \delta B^2 \rangle = \sum_i B_i^2 \langle n_i \rangle = \langle N_B \rangle \quad (14)$$

whereas for a parton system one obtains

$$\langle \delta B^2 \rangle = \frac{1}{9} \langle N_q \rangle . \quad (15)$$

Thus the ratio $\langle \delta B^2 \rangle / \langle N_{ch} \rangle \approx 0.1$ for the QGP while it is expected to be about three times larger in the coalescence model. Again, the corrections for resonance decays must be included to obtain a realistic estimate.

(iii) The constituent quark coalescence model discussed here can also be interpreted as a quark-gluon plasma far from equilibrium. This poses an apparent problem, because it has been shown in numerous papers [16] that the equilibrium scenario agrees very well with the data, in particular with those on particle ratios (chemical equilibrium). As I have already mentioned, however, particle ratios were analysed in detail by the Budapest group [9, 12] and the coalescence model was also found to be in excellent agreement with

data. I thus feel that it deserves a serious attention. Clearly, however, more work is needed to clarify fully the situation.

Acknowledgements

I would like to thank K.Fialkowski, M.Jezabek, V.Koch and M.Nowak for discussions. This investigation was supported in part by Subsydium of Foundation for Polish Science NP 1/99.

References

- [1] M.Asakawa, U.Heinz and B.Mueller, Phys.Rev.Letters 85 (2000) 2072.
- [2] S.Jeon and V.Koch, Phys.Rev.Lett. 85 (2000) 2076.
- [3] M.Bleicher, S.Jeon and V.Koch, Phys.Rev. C62 (2000) 061902; V.Koch,M.Bleicher and S.Jeon, nucl-th/0103084.
- [4] M.Asakawa, U.Heinz and B.Mueller, nucl-th/0106046.
- [5] S.Bass, P.DDanielewicz nd S.Pratt, Phys.Rev.Letters 85 (2000) 2689; S.Jeon and S.Pratt, hep-ph/0110043.
- [6] CERES coll., J.Stachel, contribution to 2002 Epiphany Conf., Cracow, January 2002 (to be published in Acta Phys. Pol. B).
- [7] C.Blume, NA49 coll., Proceedings of QM2001
- [8] J.G.Reid et al., STAR coll., Nucl.Phys. A698 (2002) 611c.
- [9] T.S.Biro, P.Levai and J.Zimanyi, Phys.Letters B347 (1995) 6; J.Zimanyi, T.S.Biro and P.Levai, J.Phys. G 23 (1997) 1941.
- [10] R.Lietava, J. Pisut, Eur.Phys. J. C5 (1998) 135; J.Pisut and N.Pisutova, Acta Phys. Pol. B28 (1997) 2817.
- [11] A.Bialas, Phys.Letters B442 (1998) 449.
- [12] J.Zimanyi et al., Phys.Letters B472 (2000) 243; T.S.Biro, P.Levai and J.Zimanyi, hep-ph/0112137 and references quoted there.

- [13] T.Csorgo, Nucl.Phys. B (Proc.Suppl.) 92 (2001) 62.
- [14] F.V.Bopp and J.Ranft, Eur.Phys.J. C22 (2001) 171; F.W.Bopp, contribution to 2002 Epiphany Conf., Cracow, January 2002 (to be published in Acta Phys. Pol. B).
- [15] M.Jezabek, private comm.
- [16] For recent results, see Proc. of the 6th Int. Conf. on Strange Quarks in Matter (Frankfurt 2001), to be published in J.Phys.G.